


/lastercode - Buchmuller et al.




#### CMSSM with large A<sub>0</sub> to get m<sub>h</sub> ~ 125 GeV

B→µ+µ- becomes an important constraint

3 parameter mSUGRA model: m<sub>3/2</sub>, m<sub>1/2</sub>, A<sub>0</sub> (tan β - fixed by B<sub>0</sub>=A<sub>0</sub>-m<sub>0</sub>) +

SubGUT universality: Min



#### Strong Moduli Stabilization/Pure Gravity Mediation

Heavy volume modulus ( $m_T \sim M_P$ ) with heavy Polonyi-like field ( $m_z >> m_{3/2}$ )


Dudas, Linde, Mambrini, Mustafayev, Olive

- The sfermion, higgsinos, heavy Higgses and gravitino have masses O(100) TeV.
- The gaugino masses are in the range of hundreds to thousands of GeV.
- The LSP is the neutral wino which is nearly degenerate with the charged wino.
- The lightest Higgs boson mass is consistent with the observed Higgs-like boson, i.e. m<sub>h</sub> ~ 125 126 GeV.

#### Strong Moduli Stabilization/Pure Gravity Mediation

#### Two parameter model!

- $m_0 = m_{3/2}; tan β$
- gaugino masses (and A-terms) generated through loops
- The sfermion, higgsinos, heavy Higgses and gravitino have masses O(100) TeV.
- The gaugino masses are in the range of hundreds to thousands of GeV.
- The LSP is the neutral wino which is nearly degenerate with the charged wino.
- The lightest Higgs boson mass is consistent with the observed Higgs-like boson, i.e. m<sub>h</sub> ~ 125 126 GeV.



#### Inflation

- as the inflaton

T - as the inflaton

$$K = -3\ln\left(T + T^* - \frac{\phi\phi^*}{3}\right)$$

$$W = M \left[ \frac{\phi^2}{2} - \frac{\phi^3}{3\sqrt{3}} \right]$$
 Ellis, Nanopolive

$$W = \sqrt{3} M \phi (T-1/2) \quad {\rm Cecotti; \, Kallosh, \, \, }$$
 Lindo

#### Equivalent to R + R<sup>2</sup> (Starobinsky) model of gravity

